

International Journal of Modern Physics A
 © World Scientific Publishing Company

Measuring $\mathcal{B}(D^+ \rightarrow \mu^+ \nu)$ and the Pseudoscalar Decay Constant f_{D^+}

N. Menea

Representing the CLEO Collaboration

201 Physics Building, Syracuse University, Syracuse, New York 13244
 E-mail: nmenaa@phy.syr.edu

In 60 pb⁻¹ of data taken on the $\psi(3770)$ resonance with the CLEO-c detector, we find 8 $D^+ \rightarrow \mu^+ \nu$ event candidates that are mostly signal, containing only 1 estimated background. Using this statistically compelling sample, we measure the value of $\mathcal{B}(D^+ \rightarrow \mu^+ \nu) = (3.5 \pm 1.4 \pm 0.6) \times 10^{-4}$, and determine $f_{D^+} = (198 \pm 41 \pm 17)$ MeV.

1. Introduction

Measuring purely leptonic decays of heavy mesons allows the determination of meson decay constants, which connect measured quantities, such as the $B\bar{B}$ mixing ratio, to CKM matrix elements. Currently, it is not possible to determine f_B experimentally from leptonic B decays, so theoretical calculations of f_B must be used.

Measurements of pseudoscalar decay constants such as f_{D^+} provide checks on these calculations and help discriminate among different models.

The decay rate is given by¹ $\Gamma(D^+ \rightarrow l^+ \nu) = \frac{G_F^2}{8\pi} f_{D^+}^2 m_l^2 M_{D^+} \left(1 - \frac{m_l^2}{M_{D^+}^2}\right)^2 |V_{cd}|^2$, where M_{D^+} is the D^+ mass, m_l is the mass of the final state lepton, V_{cd} is a CKM matrix element equal to 0.224,² and G_F is the Fermi coupling constant.

2. Analysis Technique and Event Selection

In this study we use 60 pb⁻¹ of CLEO-c data recorded at the ψ'' resonance (3.770 GeV) with the CLEO-c detector³ at the Cornell Electron Storage Ring (CESR). We fully reconstruct one charged D meson of the produced $D^+ D^-$ pairs as a tag. Tagging modes are fully reconstructed by first evaluating the difference in the energy of the decay products with the beam energy ΔE . We then require the absolute value of this difference to be within 0.02 GeV of zero, approximately twice the r.m.s. width, and then look at the reconstructed D^- beam constrained mass defined as $m_D = \sqrt{E_{beam}^2 - (\sum_i \vec{p}_i)^2}$, where i runs over all the final state particles. We then search for $D^+ \rightarrow \mu^+ \nu_\mu$ candidates with charge opposite to the charge of the tagged D^- . The charge-conjugate modes are always used.

The m_D distributions for all D^- tagging modes considered in this data sample are shown in Fig. 1. We have a total of 28574±207 tags.


2 *N. Menea*


Fig. 1. Beam Constrained Mass distributions for different fully reconstructed D^- decay candidates; the curves show the sum of Gaussian signal functions and 3^{rd} order polynomial background functions. A single signal Gaussian is used for all modes except for (a) and (c) where two Gaussians are used. (a) $D^- \rightarrow K^+ \pi^- \pi^-$, (b) $D^- \rightarrow K^+ \pi^- \pi^- \pi^0$, (c) $D^- \rightarrow K_s \pi^-$, (d) $D^- \rightarrow K_s \pi^- \pi^- \pi^+$ and (e) $D^- \rightarrow K_s \pi^- \pi^0$.

To select $D^+ \rightarrow \mu^+ \nu_\mu$ events we first reconstruct D^- event candidates and then search for events with a single additional charged track presumed to be a μ^+ . Then we infer the existence of the neutrino by requiring a measured value near zero (the neutrino mass) of the missing mass squared (MM^2) defined as $MM^2 = (E_{beam} - E_{\mu^+})^2 - (-\vec{p}_{D^-} - \vec{p}_{\mu^+})^2$ where \vec{p}_{D^-} is the three-momentum of the fully reconstructed D^- . We reject events with extra charged tracks besides the muon candidate and maximum extra shower energy above 250 MeV.

3. Results

The MM^2 distribution for our tagged events is shown in Fig. 2.⁴ We see a signal near zero containing 8 events within a $\pm 2\sigma$ interval, -0.056 GeV^2 to $+0.056 \text{ GeV}^2$. This signal is due to the $D^+ \rightarrow \mu^+ \nu_\mu$ mode we are seeking. The large peak centered




Fig. 2. MM^2 using D^- tags and one additional opposite sign charged track and no extra energetic showers (see text). The insert shows the signal region for $D^+ \rightarrow \mu^+\nu$ enlarged, $\pm 2\sigma$ range is shown between the two arrows.

near 0.25 GeV^2 is from the decay $D^+ \rightarrow \bar{K}^0 \pi^+$ that is far from our signal region.

Backgrounds from $D^+ \rightarrow \tau^+\nu$, $\tau^+ \rightarrow \pi^+\nu$, $D^+ \rightarrow \pi^+\pi^0$ and other smaller sources are estimated as summing to 1.07 events. Subtracting the background from our 8 events in the signal region, we determine a branching fraction using a detection efficiency for the single muon of 69.9% and 28574 \pm 207 D^\mp tags. We find $\mathcal{B}(D^+ \rightarrow \mu^+\nu_\mu) = (3.5 \pm 1.4 \pm 0.6) \times 10^{-4}$ and the decay constant is $f_{D^+} = (198 \pm 41 \pm 17) \text{ MeV}$.

I gratefully acknowledge my advisor Professor S. Stone for all his guidance and help.

References

1. J. L. Rosner, in **Particles and Fields 3**, Proceed. of the 1988 Banff Summer Inst., Banff, Alberta, Canada, ed. by A. N. Kamal and F. C. Khanna, World Scientific, Singapore, 1989, 395.
2. S. Eidelman *et al.* (PDG), Phys. Lett. B **592**, 1 (2004).
3. CLEO Collaboration, CLNS-01/1742 and references therein.
4. D. Besson *et al.*, "Measurement of $\mathcal{B}(D^+ \rightarrow \mu^+\nu)$ and the Pseudoscalar Decay Constant f_{D^+} ", [hep-ex/[0408071].