

Electroweak phase transition in the MSSM with four generations

S. W. Ham⁽¹⁾, S. K. Oh^(1,2), D. Son⁽¹⁾,

⁽¹⁾ *Center for High Energy Physics, Kyungpook National University
Daegu 702-701, Korea*

⁽²⁾ *Department of Physics, Konkuk University, Seoul 143-701, Korea*

Abstract

By assuming the existence of the sequential fourth generation to the minimal supersymmetric standard model (MSSM), we study the possibility of a strongly first-order electroweak phase transition. We find that there is a parameter region of the MSSM where the electroweak phase transition is strongly first order. In that parameter region, the mass of the lighter scalar Higgs boson is calculated to be above the experimental lower bound, and the scalar quarks of the third and the fourth generations are heavier than the corresponding quarks.

I. Introduction

As a mechanism to explain the baryon asymmetry of the universe, the electroweak baryogenesis is given wide attentions, since it can be tested in the future high energy experiments [1]. Sakharov have several decades ago established the three essential conditions for generating dynamically the baryon asymmetry of the universe from a baryon-symmetric universe [2]. As is well known, the three conditions are the presence of baryon number violation, the violation of both C and CP, and a departure from thermal equilibrium. The possibility of the electroweak phase transition has already been exhaustively studied, which can provide the baryon number violation and the violation of both C and CP. The remaining Sakharov condition, the departure from thermal equilibrium, may be fulfilled at a weak scale temperatures if the nature of the electroweak phase transition is first order. The difficulty of the standard model (SM) is that the strength of the first order electroweak phase transition, which must be strong enough for preserving the generated baryon asymmetry at the electroweak scale, appears too weak for the experimentally allowed mass of the SM scalar Higgs boson [3].

Thus, it seems that electroweak baryogenesis requires a new physics beyond the SM at weak scale [4]. The minimal supersymmetric standard model (MSSM) has been studied intensively within the context of electroweak baryogenesis. It is observed that, if one of the scalar top quark has a mass smaller than top quark mass, the MSSM may possess a parameter region where the electroweak phase transition is strong enough [5]. In this scenario of a light scalar top quark, the requirement that the electroweak phase transition should be strongly first order is equivalent to the imposition of an upper bound of about 120 GeV on the lightest Higgs scalar boson mass of the MSSM.

In this paper, we examine yet another possibility for the strongly first-order electroweak phase transition in the MSSM. We study the effect of the fourth generation of quarks on the strength of the first order electroweak phase transition in the MSSM. Although the number of the SM neutrino species has already been fixed experimentally as three, the existence of the sequential fourth generation is still anticipated and searched. In principle, the SM, as well as the MSSM, can accommodate any number of generations. In various contexts, the MSSM with four generations has been studied [6-8]. For us, the fourth generation of quarks are found to enhance the strength of the first-order electroweak phase transition, while the scalar Higgs boson mass is calculated to be above the experimental lower bound, and the scalar quarks of the third generations are comparable to the supersymmetry breaking scale ($M_{\text{SUSY}} = 1 \text{ TeV}$), in a reasonably wide region of parameter space in the MSSM. In our scenario, a light scalar quark is not necessary to ensure the first order electroweak phase transition be strong; the scalar quarks of the third and the fourth generations may be heavier than the corresponding quarks.

II. Higgs potential in decoupling limit without mixing

Let us study a particular, yet reasonable as well as plausible, form of the Higgs potential in the MSSM with four generations of quarks for the electroweak phase transition. We consider only the third and the fourth generations, and assume that there is no mixing between them. The fourth generation appears simply in a repetitive manner. As is well known, there are two Higgs doublets in the Higgs sector of the MSSM, namely, $H_1^T = (H_1^0, H_1^-)$ and $H_2^T = (H_2^+, H_2^0)$. After electroweak symmetry breaking emerge five physical Higgs bosons: two neutral scalar Higgs bosons (h, H), one neutral pseudoscalar Higgs bosons (A), and a pair of charged Higgs bosons (H^\pm). We assume that the mass of h is lighter than that of H . At the tree level, the Higgs sector

of the MSSM depends on only two free parameters. We take them to be the ratio $\tan\beta = v_2/v_1$ of the two real vacuum expectation values (VEVs) v_1 of H_1^0 and v_2 of H_2^0 and m_A , the mass of A . In this paper, we assume that CP is conserved in the Higgs sector by choosing all parameters in the effective Higgs potential to be real.

In the decoupling limit, where $m_A \gg m_Z$, with fixed $\tan\beta$, only one linear combination of the two neutral scalar Higgs bosons,

$$\phi = \sqrt{2} \cos\beta \text{Re}(H_1^0) + \sqrt{2} \sin\beta \text{Re}(H_2^0) , \quad (1)$$

remains light at the electroweak scale [9]. In this limit, the tree-level Higgs potential at zero temperature can be expressed in terms of ϕ as

$$V_0(\phi, 0) = -m_0^2 \phi^2 + \frac{\lambda}{4} \phi^4 . \quad (2)$$

Since all quartic terms have gauge coupling coefficients in the MSSM, the quartic Higgs self-coupling λ is given as $\lambda = (g_1^2 + g_2^2)/4$. Note that there is an upper bound on the mass of h as $m_h \leq m_Z |\cos 2\beta|$ at the tree level in the MSSM. In this limit, the couplings of h to gauge bosons and fermions are identical to the couplings of the SM Higgs boson, which implies that one cannot distinguish phenomenologically the SM scalar Higgs boson from h [10]. Thus, one might expect that m_h has the same experimental lower bound as the SM scalar Higgs boson in the decoupling limit [11]. The current experimental lower bound on the mass of the SM scalar Higgs boson is about 114.5 GeV.

Now, at the one-loop level at zero temperature, the effective Higgs potential is given by

$$V(\phi, 0) = V_0(\phi, 0) + V_1(\phi, 0) , \quad (3)$$

where the one-loop contribution $V_1(\phi, 0)$ at zero temperature is obtained via the effective potential method as [12]

$$V_1(\phi, 0) = \sum_l \frac{n_l m_l^4(\phi)}{64\pi^2} \left[\log \left(\frac{m_l^2(\phi)}{\Lambda^2} \right) - \frac{3}{2} \right] , \quad (4)$$

where l stands for various participating particles: the gauge bosons W , Z , the third generation quarks and scalar quarks t , b , \tilde{t}_1 , \tilde{t}_2 , \tilde{b}_1 and \tilde{b}_2 , as well as the fourth generation quarks and scalar quarks t' , b' , \tilde{t}'_1 , \tilde{t}'_2 , \tilde{b}'_1 and \tilde{b}'_2 . The renormalization scale in the above one-loop effective potential is set as $\Lambda = m_Z$. The degrees of freedom for each particle are: $n_W = 6$, $n_Z = 3$, $n_t = n_b = -12$, $n_{\tilde{t}_i} = n_{\tilde{b}_i} = 6$ ($i = 1, 2$), $n_{t'} = n_{b'} = -12$, $n_{\tilde{t}'_i} = n_{\tilde{b}'_i} = 6$ ($i = 1, 2$). Their field-dependent masses are given by $m_W^2(\phi) = g_2^2 \phi^2/4$, $m_Z^2(\phi) = (g_1^2 + g_2^2) \phi^2/4$, $m_t^2(\phi) = h_t^2 \sin^2 \beta^2 \phi^2/2$, $m_b^2(\phi) = h_b^2 \cos^2 \beta^2 \phi^2/2$, $m_{t'}^2(\phi) = h_{t'}^2 \sin^2 \beta^2 \phi^2/2$, $m_{b'}^2(\phi) = h_{b'}^2 \cos^2 \beta^2 \phi^2/2$, and

$$m_{\tilde{q}_1 \tilde{q}_2}^2(\phi) = \frac{m_{\tilde{q}_L}^2(\phi) + m_{\tilde{q}_R}^2(\phi)}{2} \mp \sqrt{\left(\frac{m_{\tilde{q}_L}^2(\phi) - m_{\tilde{q}_R}^2(\phi)}{2} \right)^2 + \tilde{A}_q^2 m_q^2(\phi)} , \quad (5)$$

with $q = t, b, t', b'$. In the above expression for the scalar quark masses [13], we have

$$\begin{aligned} m_{\tilde{t}_L}^2(\phi) &= m_Q^2 + m_t^2(\phi) + \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right) \cos 2\beta m_Z^2(\phi) , \\ m_{\tilde{t}_R}^2(\phi) &= m_U^2 + m_t^2(\phi) + \frac{2}{3} \sin^2 \theta_W \cos 2\beta m_Z^2(\phi) , \\ m_{\tilde{b}_L}^2(\phi) &= m_Q^2 + m_b^2(\phi) + \left(-\frac{1}{2} + \frac{1}{3} \sin^2 \theta_W \right) \cos 2\beta m_Z^2(\phi) , \end{aligned}$$

$$m_{b_R}^2(\phi) = m_D^2 + m_b^2(\phi) - \frac{1}{3} \sin^2 \theta_W \cos 2\beta m_Z^2(\phi), \quad (6)$$

and similarly for the fourth generation by substituting with primed quantities, where $\sin \theta_W$ is the weak mixing angle.

The parameters \tilde{A}_t , \tilde{A}_b , $\tilde{A}_{t'}$ and $\tilde{A}_{b'}$ in the above expressions for the scalar quark masses are given as $\tilde{A}_t = A_t - \mu \cot \beta$, and $\tilde{A}_b = A_b - \mu \tan \beta$, and similarly for the fourth generation. Note that $\tilde{A}_t = 0$ ($\tilde{A}_b = 0$) does not necessarily imply that the right-handed and the left-handed scalar top (bottom) quarks are degenerate in mass, since there is D -term contributions. Only if D -term contributions to the scalar top (bottom) quark masses are neglected, $\tilde{A}_t = 0$ ($\tilde{A}_b = 0$) would yield degenerate right-handed and left-handed scalar top (bottom) quarks.

However, we remark that the parameters \tilde{A}_t , \tilde{A}_b , $\tilde{A}_{t'}$ and $\tilde{A}_{b'}$ control the mixings between the scalar top or scalar bottom masses in each generation. If these parameters are zero, there would be no mixing between right-handed and left-handed scalar quarks of each generation. In this paper, we assume that there is no mixing, taking $\tilde{A}_t = \tilde{A}_b = \tilde{A}_{t'} = \tilde{A}_{b'} = 0$ in the expressions for the scalar quark masses. Therefore, we study the MSSM Higgs potential in the decoupling limit without mixing.

The decoupling limit without mixing is an optimal situation for electroweak phase transition to be strongly first order. The case without mixing is more favorable for the first order electroweak phase transition to be strong than the case with mixing, as the strength of the electroweak phase transition is found to decrease when the mixings between scalar quarks are taken into account [13]. Also, one can notice, for example, in Fig. 2 of Ref. [14], that the strength of the electroweak phase transition increases as m_A increases in the MSSM with three generations. Thus, we study the MSSM Higgs potential in circumstances that enhance the first order electroweak phase transition.

Now, the renormalized parameter m_0^2 in the Higgs potential can be eliminated by the minimum condition. By calculating the first derivative of $V(\phi, 0)$ with respect to ϕ , m_0^2 is expressed as

$$\begin{aligned} m_0^2 = & \frac{1}{2} m_Z^2 \cos^2 2\beta + \frac{3m_W^4}{8\pi^2 v^2} \left[\log \left(\frac{m_W^2}{\Lambda^2} \right) - 1 \right] + \frac{3m_Z^4}{16\pi^2 v^2} \left[\log \left(\frac{m_Z^2}{\Lambda^2} \right) - 1 \right] \\ & - \frac{3m_t^4}{4\pi^2 v^2} \left[\log \left(\frac{m_t^2}{\Lambda^2} \right) - 1 \right] - \frac{3m_b^4}{4\pi^2 v^2} \left[\log \left(\frac{m_b^2}{\Lambda^2} \right) - 1 \right] \\ & - \frac{3m_{t'}^4}{4\pi^2 v^2} \left[\log \left(\frac{m_{t'}^2}{\Lambda^2} \right) - 1 \right] - \frac{3m_{b'}^4}{4\pi^2 v^2} \left[\log \left(\frac{m_{b'}^2}{\Lambda^2} \right) - 1 \right] \\ & + \sum_a \frac{3m_a^2}{8\pi^2 v^2} (m_a^2 - m_Q^2) \left[\log \left(\frac{m_a^2}{\Lambda^2} \right) - 1 \right] \\ & + \sum_{a'} \frac{3m_{a'}^2}{8\pi^2 v^2} (m_{a'}^2 - m_{Q'}^2) \left[\log \left(\frac{m_{a'}^2}{\Lambda^2} \right) - 1 \right], \end{aligned} \quad (7)$$

where $a = \tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$, and $a' = \tilde{t}'_1, \tilde{t}'_2, \tilde{b}'_1, \tilde{b}'_2$, and $v = 246$ GeV. The mass of h at the one-loop level in the decoupling limit without mixing at zero temperature is obtained by calculating the second derivative of $V(\phi, 0)$ with respect to ϕ as

$$\begin{aligned} m_h^2 = & m_Z^2 \cos^2 2\beta + \frac{3m_W^4}{4\pi^2 v^2} \log \left(\frac{m_W^2}{\Lambda^2} \right) + \frac{3m_Z^4}{8\pi^2 v^2} \log \left(\frac{m_Z^2}{\Lambda^2} \right) \\ & - \frac{3m_t^4}{2\pi^2 v^2} \log \left(\frac{m_t^2}{\Lambda^2} \right) - \frac{3m_b^4}{2\pi^2 v^2} \log \left(\frac{m_b^2}{\Lambda^2} \right) \end{aligned}$$

$$\begin{aligned}
& -\frac{3m_{t'}^4}{2\pi^2v^2} \log\left(\frac{m_{t'}^2}{\Lambda^2}\right) - \frac{3m_{b'}^4}{2\pi^2v^2} \log\left(\frac{m_{b'}^2}{\Lambda^2}\right) \\
& + \sum_a \frac{3(m_a^2 - m_Q^2)^2}{4\pi^2v^2} \log\left(\frac{m_a^2}{\Lambda^2}\right) + \sum_{a'} \frac{3(m_{a'}^2 - m_{Q'}^2)^2}{4\pi^2v^2} \log\left(\frac{m_{a'}^2}{\Lambda^2}\right), \tag{8}
\end{aligned}$$

where $a = \tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2$, and $a' = \tilde{t}'_1, \tilde{t}'_2, \tilde{b}'_1, \tilde{b}'_2$.

Now, we study the effect of finite temperature. The one-loop contribution at finite temperature is given by [15]

$$V_1(\phi, T) = \sum_l \frac{n_l T^4}{2\pi^2} \int_0^\infty dx x^2 \log \left[1 \pm \exp \left(-\sqrt{x^2 + m_l^2(\phi)/T^2} \right) \right], \tag{9}$$

where $l = W, Z, t, b, \tilde{t}_1, \tilde{t}_2, \tilde{b}_1, \tilde{b}_2, t', b', \tilde{t}'_1, \tilde{t}'_2, \tilde{b}'_1, \tilde{b}'_2$. and the negative sign is for bosons and the positive sign for fermions. The full one-loop effective potential at finite temperature that we are considering can now be expressed as

$$V(\phi, T) = V_0(\phi, 0) + V_1(\phi, 0) + V_1(\phi, T). \tag{10}$$

We perform the exact integration in $V(\phi, T)$ instead of employing the high-temperature approximation.

III. Numerical Analysis

At the tree level, we have only one free parameter $\tan\beta$ in the decoupling limit where $m_A \gg m_Z$. At the one-loop level, the number of free parameters increases as $m_t, m_b, m_{t'}, m_{b'}$, and $m_Q, m_{Q'}$ are introduced to the one-loop contributions. To be concrete, we set $m_t = 175$ GeV and $m_b = 4.5$ GeV, and we take for simplicity the soft SUSY breaking parameters as $m_Q^2 = m_{Q'}^2 = m_D^2$, and similarly for the fourth generation. Since we assume no mixing in the scalar quark sector, we set $\tilde{A}_t = \tilde{A}_b = \tilde{A}_{t'} = \tilde{A}_{b'} = 0$. For the masses of the fourth generation quarks, there are some experimental constraints. Some years ago, Tevatron data have set $m_{b'} > 119$ GeV, and recently, from the search for long-lived charged massive particles at Tevatron come more stringent experimental lower bounds of $m_{b'} > 180$ GeV and $m_{t'} > 230$ GeV [16]. With these constraints in mind, we take $m_{t'} = 250$ GeV and $m_{b'} = 200$ GeV. Finally, we take 1 TeV for the value of m_Q from the SUSY breaking scale $M_{\text{SUSY}} = 1$ TeV. Thus, our numerical analysis involves two free parameters: $\tan\beta$ and $m_{Q'}$.

In Fig. 1, we show a typical behavior of $V(\phi, T)$ as a function of ϕ , at a critical temperature $T = T_c = 84.225$ GeV, where we take $\tan\beta = 20$ and $m_{Q'} = 100$ GeV. As one can see in the figure, we obtain the critical vacuum expectation value as $v_c = 149$ GeV, and the ratio as $v_c/T_c = 1.769$. one can notice that the potential in Fig. 1 allows a strongly first-order electroweak phase transition. The scalar quark masses are obtained as $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, $m_{\tilde{b}_2} = 1001$ GeV, $m_{\tilde{t}'_1} = 263$ GeV, $m_{\tilde{t}'_2} = 266$ GeV, $m_{\tilde{b}'_1} = 225$ GeV, and $m_{\tilde{b}'_2} = 231$ GeV.

For the mass of h , we obtain $m_h = 129$ GeV with the parameter values of Fig. 1. This number is a little larger than the experimental lower bound on the SM Higgs boson mass, 115 GeV. In the decoupling limit of $m_A \gg m_Z$, the behavior of h is identical to that of the SM Higgs boson. The search for a light Higgs boson in the MSSM by DELPHI collaboration without mixing, where only three generations of quarks are taken into account, suggests that m_h is about

115 GeV, for $\tan\beta > 15$ [17]. Comparing this number with our result of 129 GeV in Fig. 1, we may well deduce that the difference is due to the contribution by the fourth generation of quarks. The contribution by the fourth generation of quarks also enables the electroweak phase transition strongly first order. Our value is compatible with the result of Ref. [6] where the upper bound on the lightest Higgs boson mass is obtained as 152 GeV for small $\tan\beta$ and $96 \leq m_{t'}, m_{b'} \leq 125$ GeV in the MSSM with four generations of quarks. If the mass difference between t' and b' is small, they may have smaller masses [7].

Now, let us study other regions of the parameter space. We vary $m_{Q'}$ while fixing $\tan\beta = 20$. The masses of the third generation of scalar quarks are then the same as Fig. 1: $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, and $m_{\tilde{b}_2} = 1001$ GeV. The masses of the fourth generation of scalar quarks, as well as other relevant quantities, are shown in Table I. The second row of Table I for $m_{Q'} = 100$ GeV corresponds to the numerical result of Fig. 1. One can see that the strength of the first order electroweak phase transition decreases as $m_{Q'}$ increases until $m_{Q'}$ reaches 140 GeV, beyond which v_c/T_c becomes less than 1. Thus, the electroweak phase transition remains strongly first order for $m_{Q'} \leq 140$ GeV.

We need to explore the boundary of the parameter space beyond which the electroweak phase transition is no longer strongly first order. In Table I, one can see that $v_c/T_c \sim 1$ for $\tan\beta = 20$ and $m_{Q'} = 140$ GeV. From this point, we examine several points of $(m_{Q'}, \tan\beta)$ which yield $v_c/T_c \sim 1$, by adjusting T_c . For each value of $\tan\beta$, we find the upper bound value of $m_{Q'}$, beyond which v_c/T_c becomes less than 1. The result is shown in Table II. The fourth row of Table II is corresponds to the fourth row of Table I. The numbers in Table II indicate that the electroweak phase transition can be strongly first order for $2 \leq \tan\beta \leq 40$ if $m_{Q'} \leq 140$ GeV.

In Fig. 2, we plot the numerical results of Table 2 on $(m_{Q'}, \tan\beta)$ -plane. The dashed curves denote the contours of m_h . The solid curve denotes the contour of $v_c/T_c = 1$. On the left-hand side of the solid curve we have $v_c/T_c \geq 1$, that is, the electroweak phase transition is strongly first order in the region to the left of the solid curve. Consequently, we find a region in the parameter space of the MSSM with four generations of quarks where a strongly first order electroweak phase transition is allowed. The parameter values of the allowed region are within the experimental constraints, and yield m_h consistent with experimental lower bound.

Table I: Some values of $m_{Q'}$ for which the first order electroweak phase transition is strong ($v_c/T_c > 1.0$). The critical temperatures T_c are obtained for which the finite temperature effective potential has two degenerate vacua. The relevant parameter values are the same as Fig. 1, that is, $\tan\beta = 20$, $m_Q = 1$ TeV, $m_{t'} = 250$ GeV, and $m_{b'} = 200$ GeV.

$m_{Q'}$	$m_{\tilde{t}_1}$	$m_{\tilde{t}_2}$	$m_{\tilde{b}_1}$	$m_{\tilde{b}_2}$	m_h	T_c	v_c	v_c/T_c
50	249	252	207	214	122	76.090	185	2.431
100	263	266	225	231	129	84.225	149	1.769
130	276	279	239	245	134	90.180	113	1.253
140	281	284	245	251	136	92.268	93	1.007
150	286	289	251	256	138	94.384	76	0.805

Table II: Some values of $(m_{Q'}, \tan \beta)$ beyond which the first order electroweak phase transition is strong ($v_c/T_c \sim 1.0$). The relevant parameters values are the same as Table I. The masses of the scalar quarks of the fourth generation are calculated as about $m_{\tilde{t}_1} = 282$ GeV, $m_{\tilde{t}_2} = 285$ GeV, $m_{\tilde{b}_1} = 246$ GeV, $m_{\tilde{b}_2} = 251$ GeV.

$\tan \beta$	$m_{Q'}$	m_h	T_c	v_c	v_c/T_c
2	147	120	87.163	88	1.009
5	142	133	91.256	92	1.008
10	140	135	91.965	92	1.000
20	140	136	92.268	93	1.007
30	140	137	92.326	93	1.007
40	140	140	92.346	93	1.007

IV. Conclusions

Up to now, we study the possibility of a strongly first order electroweak phase transition in the MSSM with sequential four generations of quarks. We assume that $m_A \gg m_Z$ and $\tilde{A}_l = 0$ ($l = t, b, t', b'$), that is, we work in the decoupling limit and in the case of no mixing between scalar quarks. We choose the relevant parameter values to be: $m_t = 175$ GeV, $m_b = 4.5$ GeV, $m_{t'} = 250$ GeV, $m_{b'} = 200$ GeV, and $m_Q = 1$ TeV. We take $m_Q^2 = m_U^2 = m_D^2$, and similarly for the fourth generation. These numbers are consistent with experimental constraints.

We search the parameter space of $(m_{Q'}, \tan \beta)$ -plane to examine if the electroweak phase transition is strongly first order. We find that there are regions in the $(m_{Q'}, \tan \beta)$ -plane that satisfy our criterion of $v_c/T_c \geq 1$. For $2 \leq \tan \beta \leq 40$, the electroweak phase transition is strongly first order in the region where $m_{Q'} \leq 140$ GeV. The scalar quark masses of the fourth generation are controlled mainly by the soft SUSY breaking parameter $m_{Q'}$. In the region where the electroweak phase transition is strongly first order, the scalar quark masses of the fourth generation are obtained to be larger than the quark masses of the fourth generation.

Acknowledgments

This work was supported by Korea Research Foundation Grant (2001-050-D00005).

Reference

- [1] V.A. Kuzmin, V.A. Rubakov, and M.E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985); M.E. Shaposhnikov, JETP Lett. **44**, 465 (1986); Nucl. Phys. B **287**, 757 (1987); Nucl. Phys. B **299**, 797 (1988); L. McLerran, Phys. Rev. Lett. **62**, 1075 (1989); N. Turok and J. Zadrozny, Phys. Rev. Lett. **65**, 2331 (1990); Nucl. Phys. B **358**, 471 (1991); L.D. McLerran, M.E. Shaposhnikov, N. Turok, and M.B. Voloshin, Phys. Lett. B **256**, 451 (1991); M. Dine, P. Huet, R.J. Singleton, and L. Susskind, Phys. Lett. B **257**, 351 (1991); A.G. Cohen, D.B. Kaplan, and A.E. Nelson, Annu. Rev. Nucl. Part. Sci. **43**, 27 (1993); M. Trodden, Rev.

- Mod. Phys. **71**, 1463 (1999); A. Riotto and M. Trodden, Annu. Rev. Nucl. Part. Sci. **49**, 35 (1999).
- [2] A.D. Sakharov, JETP Lett. **5**, 24 (1967).
- [3] A.I. Bochkarev, S.V. Kuzmin, and M.E. Shaposhnikov, Mod. Phys. Lett. A **2**, 417 (1987); Phys. Lett. B **244**, 275 (1990); M. Dine, R.G. Leigh, P. Huet, A. Linde, and D. Linde, Phys. Rev. D **46**, 550 (1992); P. Arnold and O. Espinosa, Phys. Rev. D **47**, 3546 (1993); Z. Fodor and A. Hebecker, Nucl. Phys. B **432**, 127 (1994); K. Kajantie, M. Laine, K. Rummukainen, and M. Shaposhnikov, Phys. Rev. Lett. **77**, 2887 (1996); F. Csikor, Z. Fodor, and J. Heitger, Phys. Rev. Lett. **82**, 21 (1999).
- [4] A.I. Bochkarev, S.V. Kuzmin and M.E. Shaposhnikov, Phys. Rev. D **43**, 369 (1991); G.W. Anderson and L.J. Hall, Phys. Rev. D **45**, 2685 (1992); N. Turok and J. Zadrozny, Nucl. Phys. B **369**, 729 (1992); J. Choi and R.R. Volkas, Phys. Lett. B **317**, 385 (1993); K. Enqvist, K. Kainulainen, and I. Vilja, Nucl. Phys. B **403**, 749 (1993); M. Pietroni, Nucl. Phys. B **402**, 27 (1993); I. Vilja, Phys. Lett. B **324**, 197 (1994); A.T. Davies, C.D. Froggatt, G. Jenkins, and R.G. Moorhouse, Phys. Lett. B **336**, 464 (1994); J. Choi, Phys. Lett. B **345**, 253 (1995); A.T. Davies, C.D. Froggatt, and R.G. Moorhouse, Phys. Lett. B **372**, 88 (1996); J.M. Cline and P.A. Lemieux, Phys. Rev. D **55**, 3873 (1997); S.J. Huber and M.G. Schmidt, Eur. Phys. J. C **10**, 473 (1999); M. Bastero-Gil, C. Hugonie, S.F. King, D.P. Roy, and S. Vempati, Phys. Lett. B **489**, 359 (2000); A. Menon, D.E. Morrissey, and C.E.M. Wagner, Phys. Rev. D **70**, 035005 (2004); S.W. Ham, S.K. Oh, C.M. Kim, E.J. Yoo, and D. Son, Phys. Rev. D **70**, 075001 (2004); J. Kang, P. Langacker, T. Li, and T. Liu, hep-ph/0402086; M. Carena, A. Megevand, M. Quirós, and C.E.M. Wagner, hep-ph/0410352.
- [5] M. Carena, M. Quirós and C.E.M. Wagner, Phys. Lett. B **380**, 81 (1996); D. Delepine, J.M. Gerard, R.G. Felipe, and J. Weyers, Phys. Lett. B **386**, 183 (1996); M. Carena, M. Quirós and C.E.M. Wagner, Nucl. Phys. B **524**, 3 (1998); M. Laine and K. Rummukainen, Phys. Rev. Lett. **80**, 5259 (1998); Nucl. Phys. B **535**, 423 (1998); J.M. Cline and G.D. Moore, Phys. Rev. Lett. **81**, 3315 (1998); K. Funakubo, A. Kakuto, S. Otsuki, and F. Toyoda, Prog. Theor. Phys. **99**, 1045 (1998); K. Funakubo, Prog. Theor. Phys. **101**, 415 (1999); M. Losada, Nucl. Phys. B **537**, 3 (1999).
- [6] J.E. Dubicki and C.D. Froggatt, Phys. Lett. B **567**, 46 (2003).
- [7] J.F. Gunion, D.W. McKay, and H. Pois, Phys. Rev. D **53**, 1616 (1996).
- [8] M. Carena, H.E. Haber, and C.E.M. Wagner, Nucl. Phys. B **472**, 55 (1996).
- [9] G.F. Giudice, Phys. Rev. D **45**, 3177 (1992).
- [10] M. Carena and H.E. Haber, Prog. Part. Nucl. Phys. **50** 63 (2003).
- [11] C. Balazs, M. Carena, and C.E.M. Wagner, Phys. Rev. D **70**, 015007 (2004).
- [12] S. Coleman and E. Weinberg, Phys. Rev. D **7**, 1888 (1973).
- [13] J.R. Espinosa, M. Quirós, and F. Zwirner, Phys. Lett. B **307**, 106 (1993).
- [14] B. de Carlos and J.R. Espinosa, Nucl. Phys. B **503**, 24 (1997).

- [15] L. Dolan and R. Jackiw, Phys. Rev. D **9**, 3320 (1974).
- [16] CDF Collaboration, Phys. Rev. Lett. **84**, 835 (2000); Phys. Rev. Lett. **90**, 131801.
- [17] DELPHI Collaboration, Eur. Phys. J. C **32**, 145 (2004).

Figure Captions

Fig. 1: The plot of $V(\phi, T)$ as a function of ϕ , for the critical temperature $T_c = 84.225$ GeV. The relevant parameter values are set as follows: $\tan\beta = 20$, $m_Q = 1$ TeV, $m_{t'} = 250$ GeV, $m_{b'} = 200$ GeV, and $m_{Q'} = 100$ GeV. One can see that the electroweak phase transition is first order. The critical VEV is obtained as $v_c = 149$ GeV. Thus, $v_c/T_c = 1.769$, and the first order electroweak phase transition is strong. The potential in this figure yields $m_h = 129$ GeV, $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, $m_{\tilde{b}_2} = 1001$ GeV, $m_{\tilde{\tau}_1} = 263$ GeV, $m_{\tilde{\tau}_2} = 266$ GeV, $m_{\tilde{\nu}_1} = 225$ GeV, and $m_{\tilde{\nu}_2} = 231$ GeV.

Fig. 2: Plots of the lightest scalar Higgs boson mass and the criterion of the strongly first order electroweak phase transition in the $(m_{Q'}, \tan\beta)$ -plane. The remaining relevant parameters are set as $m_Q = 1$ TeV, $m_{t'} = 250$ GeV, and $m_{b'} = 200$ GeV. The solid curve is the contour of $v_c/T_c = 1$, and the dashed curves are the contours of $m_h = 115, 120, 125, 130, 135,$ and 140 GeV. The masses of the scalar quarks of the third generation are calculated the same as Fig. 1, that is, $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, and $m_{\tilde{b}_2} = 1001$ GeV. The region to the left-hand side of the solid curve and above the dashed curve of $m_h = 115$ GeV is where the first order electroweak phase transition is strong ($v_c/T_c \geq 1$) and m_h is consistent with the experimental lower bound.




Fig. 1: The plot of $V(\phi, T)$ as a function of ϕ , for the critical temperature $T_c = 84.225$ GeV. The relevant parameter values are set as follows: $\tan \beta = 20$, $m_Q = 1$ TeV, $m_{t'} = 250$ GeV, $m_{b'} = 200$ GeV, and $m_{Q'} = 100$ GeV. One can see that the electroweak phase transition is first order. The critical VEV is obtained as $v_c = 149$ GeV. Thus, $v_c/T_c = 1.769$, and the first order electroweak phase transition is strong. The potential in this figure yields $m_h = 129$ GeV, $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, $m_{\tilde{b}_2} = 1001$ GeV, $m_{\tilde{\nu}_1} = 263$ GeV, $m_{\tilde{\nu}_2} = 266$ GeV, $m_{\tilde{\nu}'_1} = 225$ GeV, and $m_{\tilde{\nu}'_2} = 231$ GeV.




Fig. 2: Plots of the lightest scalar Higgs boson mass and the criterion of the strongly first order electroweak phase transition in the $(m_{Q'}, \tan \beta)$ -plane. The remaining relevant parameters are set as $m_Q = 1$ TeV, $m_{t'} = 250$ GeV, and $m_{b'} = 200$ GeV. The solid curve is the contour of $v_c/T_c = 1$, and the dashed curves are the contours of $m_h = 115, 120, 125, 130, 135,$ and 140 GeV. The masses of the scalar quarks of the third generation are calculated the same as Fig. 1, that is, $m_{\tilde{t}_1} = 1013$ GeV, $m_{\tilde{t}_2} = 1014$ GeV, $m_{\tilde{b}_1} = 1000$ GeV, and $m_{\tilde{b}_2} = 1001$ GeV. The region to the left-hand side of the solid curve and above the dashed curve of $m_h = 115$ GeV is where the first order electroweak phase transition is strong ($v_c/T_c \geq 1$) and m_h is consistent with the experimental lower bound.